

Pinnacle Pediatrics

Scott R. Serbin, M.D., F.A.A.P.

Pinnacle Pediatrics Newsletter

Volume XIV, #1

Our first Newsletter of the year is always our “What to do when your child is sick” issue. The timing is appropriate. This is the time of year when most illnesses flourish – colds, coughs, sore throats, Flu, gastroenteritis, etc. I do encourage you to read this thoroughly, and to keep it in a handy location, for instant reference if your child is ill (it will be posted on our website, pinnaclepediatrics.com, under Newsletters, as well). This does thoroughly explain my advice on diagnosis and treatment of the most common illnesses you are likely to see in your children. There are always changes and additions to this advice, as medicine is an ever-evolving field of science, so please do review this even if you have read it previously.

A frequent question pertaining to childhood illness is when to keep a child home from school. From the child’s perspective, if they are feeling well enough to learn and sit in a classroom, I encourage you to send them. It will not make them sicker, or decrease their ability to fight the infection, by sitting in a classroom. I know many of you disagree with me, but I am fine sending a child with a cold to school (unless they are unusually ill with their cold). I do believe there is value in learning to function under less than ideal conditions, such as when you are mildly ill. (I have never missed a day of work due to illness or injury, and that includes Mono and several broken bones, as well as many minor illnesses bequeathed to me by my patients. I do not mean to brag, and realize I am fortunate to have avoided serious illness, but, I do value fortitude). Children CAN go to school with minor illnesses.

There is a case to be made that ill children should stay home so as not to spread their germs to others. Many schools require students to be fever-free for at least 24 hours before returning to school, and schools have various other rules regarding illness as well. That is their prerogative. However, unless you home-school your children and rarely let them venture out in public, they are going to be exposed to germs, and most people are comfortable with that.

So, if your child does encounter one of these malevolent microorganisms, read on...

Fever

Almost all of you have heard me preach that fever is not dangerous, it is simply a sign of infection. My concern is not the fever, but what is causing the fever - what is the infection and, even more important, how serious does it appear to be. Any time a child has a fever, or any symptom of illness, the most important questions to ask are..."How is the child acting?" and "How is the child drinking?" If these two items seem to be okay, then it is very unlikely that there is a serious problem. Conversely, if the child is extremely irritable or lethargic, or refusing to drink for an extended period, then we need to be concerned.

Any time a child has a fever or is ill, she is entitled to act "sick", just not "real sick". She may be fussy, sleepy, not eat well. But, she needs to drink, she needs to be arousable, consolable, and interactive to reassure us that there is nothing serious going on, that she is not "toxic".

Once it has been established that the child is not toxic, then look for other "clues" as to the source of the fever, i.e. cold symptoms (runny nose, cough, congestion, sneezing), gastrointestinal symptoms (abdominal pain, vomiting, diarrhea), sore throat, earache, etc. Often in children, there are no symptoms other than fever. Most of the time, these kids have a viral illness, which may simply run its course (usually 3-5 days) without any other symptoms.

Fever itself is not dangerous unless it reaches 107°F or higher, which is rarely seen except in severe heat stroke -- almost never with an infection. It is true that about 10% of children under 7 years of age will have a seizure with fever. But this is related to the rate of rise of the fever not how high it is. Most of the time the parent doesn't even know their child has a fever before the seizure. Fortunately, although febrile seizures are frightening to the parent, they are rarely serious. It has never been demonstrated that we can prevent febrile seizures by aggressively treating the fever.

The key is not to focus on the fever. It is worthwhile to measure the temperature one time to document that there is a true fever. (Often kids feel warm to a parent's touch, but the temperature is normal. This is not a concern. There is no disease state associated with this). After that, put the thermometer away - it is not important whether the temp. is 101 or 104. The degree of the fever correlates poorly with the severity of the infection. If the child is uncomfortable with the fever (usually the case), feel free to treat the child with an antipyretic (fever reducer). Reducing the fever will not "mask" a serious illness, and if the fever is reduced, the child will likely drink better and act better, thus reassuring us that he is not "toxic". Do not be concerned, however, if the medicine does not decrease the fever - it has been clearly shown that the response to antipyretics is not indicative of the severity of the illness. Acetaminophen (Tylenol) or Ibuprofen (Motrin, Advil) are both effective at the proper dose (15mg./kg. every four hours for Acetaminophen, 10mg./kg. every six hours for Ibuprofen). Head-to-head, Ibuprofen appears to be slightly more effective than Acetaminophen. Although you will hear medical personnel recommending alternating the two medicines, I do not believe that this is a good idea. It is hard to coordinate an every four hour and an every six hour dosing, and many mistakes, leading to overdoses, have been made in this manner. Stick with one antipyretic and use it appropriately. Besides, the main point here is that Fever is Not the Enemy. We treat it

to make the child comfortable, but the real concerns are as we discussed above. (Note -- Many cold medicines contain Acetaminophen, so combining Acetaminophen with a cold product can lead to an overdose of Acetaminophen. Always read labels to avoid this serious complication). (Also note -- Acetaminophen is the most common accidental medication poisoning in the U.S. This can lead to serious, even fatal, liver injury. Keep Acetaminophen, like all medicines, safely away from small children).

This discussion does not apply to the infant under three months of age. Although fever is not dangerous for this child either, a child under three months of age with a true fever (temp. greater than 100.5°F.) has a 20% chance of having a serious infection, and thus necessitates a call to the Pediatrician. Likewise, if a child greater than three months of age appears toxic, or the fever lasts more than 3-5 days, the Pediatrician should be called.

Colds

Colds are ubiquitous - everybody gets them. Because there are numerous viruses that cause colds, a child can get many colds in the same season. The usual symptoms are low-grade fever (99°-102°F.) for the first few days, sore throat, runny nose, sneezing, congestion, and cough. The runny nose usually starts out clear, then turns cloudy around day 4, then turns clear again around day 7. A cold may make a child slightly uncomfortable, slightly lethargic and cause a decrease in appetite. But, most kids will still be fairly active and still drink well. On average children get 6-12 colds per year. Parents often become concerned that their child is getting too many colds, and question if they have a problem with their immune system. Children who have true immune deficiencies are prone to recurrent serious infections, not colds.

Cough is the single most common reason for a call to the Pediatrician. It makes the child uncomfortable, which makes the parents uncomfortable. Parents often focus on the nature of the cough (dry, wet, harsh, phlegmy, etc.), but this is usually insignificant in determining the severity of the child's illness. The only accompanying symptom that should cause concern is difficulty breathing, particularly when the child is not in the middle of a coughing spell. If the child is breathing fast or hard for a prolonged period, the Pediatrician should be notified.

Cough often persists for 4-8 weeks, which drives parents crazy. This is due to inflammation of the airway, not the infection that initiated it, so these children are not contagious. As I have discussed in prior newsletters, cold and cough medicines are largely ineffective. Due to possible side-effects, they are no longer recommended in children under 6 years of age. Therapies that may make your child a little more comfortable include moisture in the air (vaporizer/humidifier, bathroom steam) and sipping beverages/sucking on lozenges (bathing the cough receptors in the back of the throat helps to decrease cough). Honey (for the child over 1 year of age) has also been shown to be mildly beneficial for cough. Salt-water (saline) nose drops with suction can help to ease congestion in the infant. You can make saline nose drops by mixing ¼ teaspoon of salt in 4 ounces of water. Heat it so it goes into solution, then let it cool -- Voila, saline nose drops. Vicks under the nose may offer relief from congestion, though it has no apparent benefit when rubbed on the chest.

In the child above 6 years of age, cough/cold medicines are still of questionable benefit, but have decreased risk. Pseudoephedrine is the most effective oral decongestant, but is now stored only behind the pharmacist's counter because of its role as an ingredient in the production of crystal meth, so you need to ask for it. Potential adverse effects include insomnia, headache, excitability, nervousness, decreased appetite, increased heart rate and blood pressure, arrhythmias, nausea and vomiting. Phenylephrine has replaced Pseudoephedrine in most OTC cold medicines. Numerous studies show it to be no more effective than placebo (The Medical Letter, Dec. 2015). Afrin nasal spray is effective in relieving congestion, but even when limited to 2-3 days, usage may still result in a "rebound" of nasal congestion. Dextromethorphan is the most common OTC cough suppressant, but it is not very effective. Delsym is a long acting form of Dextromethorphan that may be useful for night time cough. Previously, we would prescribe Codeine for the older child with a severe cough, but this is no longer recommended due to numerous reports of respiratory depression and death secondary to this therapy. Antibiotics have no role in treating the common cold, which is due to a virus, as they only treat bacterial infections. Echinacea, Vitamin C and Zinc have all been purported to help alleviate cold symptoms, but there is no good scientific evidence that this is true in children. Grandma's chicken soup (and actually, just Grandma) may provide the most comfort. "A cold will last seven days if you treat, one week if you don't." A good review of cold remedies can be found in the January, 2018 edition of Consumer Reports.

Frequently, a parent becomes concerned that their child's upper respiratory infection is a bacterial infection. This is usually due to a change to cloudy nasal discharge (though, as discussed, this is the norm around day 4) or the length of the symptoms. Most colds do last 7-10 days, and 2 weeks is not unusual. Cough may last 4-6 weeks, which is a frequent cause of concern. The typical bacterial upper respiratory infection (sinus infection) usually presents at the tail end of a cold. Symptoms include high fever, marked congestion, a large amount of thick yellow or green nasal discharge, and a significant worsening of the child's activity level and appetite. These symptoms should prompt a call to the Pediatrician, as sinus infections are amenable to antibiotic therapy.

FLU

Influenza, or the Flu, usually presents with the rapid onset of high fever, chills, and body aches. Other symptoms include sore throat, cough and vomiting. The symptoms of the Flu usually last for 7 days. The Flu almost always presents in epidemic fashion in the winter, not episodically throughout the year.

Influenza is a viral infection, so antibiotics are not useful. There are anti-viral drugs which treat Influenza. However, the older drugs, Amantadine and Rimantadine, are no longer very effective, and the newer drugs, primarily Tamiflu and Relenza, are not extremely effective and appear to be causing an increased number of significant side-effects. Thus, we usually do not treat Influenza except in patients requiring hospitalization, but merely offer supportive therapy, i.e. fluids and antipyretics. Consequently, we usually do not test for Influenza, since the result will not change our management. Tamiflu is being heavily advertised by the manufacturer this year, but even their ads admit that use of Tamiflu decreases the duration of the illness by only one day. It is reasonable to consider treatment with Tamiflu for children with Flu-like symptoms

of less than 48 hour duration who have significant underlying health conditions, such as heart disease or asthma or those under 2 years of age. Tamiflu prophylaxis for exposure to individuals with Flu is not recommended. The most common adverse effects of Tamiflu are nausea, vomiting and headache. More worrisome have been numerous reports of neuropsychiatric events.

Unfortunately, the current Flu season (2017 – 2018) appears to be a relatively severe one, both in terms of number of people affected and severity of illness. This is often the case when the H3N2, type A strain predominates, which it is so far this year. We have not seen much Influenza in our practice so far, which leads me to believe that the vaccine may be more effective this year than forecast, as we have a very high percentage of our patients who do get the vaccine. Or, maybe our patients just haven't been exposed to all the Flu-infected kids in a Pediatric waiting room. Hopefully our trend will continue.

SORE THROAT

In general, a sore throat (pharyngitis/tonsillitis) is due to either a virus or a bacteria. The usual bacteria that causes a sore throat is Streptococcus, or "strep". Viruses are responsible for 90% of sore throats, although in "strep season", March and April, strep may cause 50% of sore throats.

Often, a cold may start out as just a sore throat, and then on day 2 or 3 the child will develop a full-blown cold. Strep throat usually presents with a high fever, severe sore throat, bright red tonsils (often with pus) and large, swollen lymph nodes in the neck. It is often associated with a headache, abdominal pain and vomiting. Occasionally, strep throat will also be accompanied by a fine, pimply, "sand-paper-like" rash - this is called "Scarlet Fever". Although many years ago this was a more serious form of strep, today it does not represent a more severe illness.

Studies done over 40 years ago demonstrated that it was difficult to distinguish between viral pharyngitis and strep throat. Consequently, physicians have relied on throat cultures and rapid strep tests to make the correct diagnosis. However, these tests are very uncomfortable for most children, and their consequent lack of cooperation often results in an unsatisfactory throat swab, yielding an invalid test. In addition, 5% of the population will have a positive strep test, despite not having an active infection. I am now convinced with 30+ years of clinical experience, that basing treatment on my clinical judgement may be a better option than doing a throat swab. Although I will continue to do rapid strep tests under certain circumstances, I will be doing fewer of them going forward and basing treatment decisions on clinical criteria. (I hear the cheers from the extreme gaggers).

So, if your child has the symptoms of strep that I described, I will likely treat with antibiotics. However, if he/she has only had a sore throat for 1-2 days (which is likely the prelude to a cold), or if your child has other viral symptoms (runny nose, congestion, sneezing, cough), this is likely a viral pharyngitis, and does not require antibiotic treatment.

There is no rush to treat a child with strep throat. Antibiotics initiated within 18 days of the onset of infection will prevent Rheumatic Fever, our chief concern with strep (although only 0.1% of cases of Strep throat result in Rheumatic Fever).

There is no treatment for a viral pharyngitis, just supportive measures such as pain relievers, Chloraseptic spray/lozenges (this contains Benzocaine, a local numbing agent – o.k. for kids over 6 years old) and fluids. Most viral sore throats last 3-5 days, though some, particularly those caused by Coxsackie virus (Hand-Foot-Mouth disease) last for 7 days.

PINK EYE

Pink eye, or conjunctivitis, is an infection of the conjunctival lining of the eye. This can be due to a virus or a bacteria. The primary way to assess the etiology (without doing a culture) is based on the amount of discharge from the eye. A viral conjunctivitis causes erythema (redness) of the inner lower eyelid and the sclera (the white part of the eyeball), but only causes minimal discharge (greater on awaking, then 3-4 times during the day). A bacterial conjunctivitis also causes erythema, but produces a large amount of discharge that accumulates constantly throughout the day.

The treatment for a viral conjunctivitis is simply warm compresses. The duration of symptoms is usually 7 days. Warm compresses are also beneficial for a bacterial conjunctivitis, especially first thing in the morning when the child's eyes are glued shut (which can be very frightening to a young child). Just let the warm washcloth soak on the eyelids for 5 minutes and the eyes will gradually open. In addition, we treat bacterial conjunctivitis with topical antibiotic drops, which will hasten the resolution of the infection (assuming you have six burly Bouncers to hold the child down while you administer the drops).

Pink eye is very contagious, which is why schools and day-cares often exclude children with pink eye. However, it is not serious or dangerous, and only mildly uncomfortable. Often, a facility will advise a parent that their child cannot come back until they are being treated, not realizing that there is no treatment for most of these kids. Many times I have argued with school nurses and administrators concerning this issue, usually successfully. I do not believe children should be excluded due to a "cold in the eye", any more than they should be excluded due to a cold. This is also the official position of the American Academy of Pediatrics. The key to preventing transmission, as with so many illnesses, is washing the hands, either with soap and water or hand sanitizers, and avoiding touching other children's eyes.

GASTROENTERITIS

This is the final common illness that I will discuss. Typically, this starts with vomiting, which, fortunately, usually lasts less than 24 hours. The advice is to wait 2 hours from the last time the child vomited, and then begin sips of clear liquids (Pedialyte in the infant, any clear liquid in the older child) every 15 minutes. This is very labor intensive, as we wish to get a lot of fluid into the child, but only a little at a time. If the child vomits again, wait another 2 hours, and then

start over. Gradually increase the volume as tolerated. If the child has a fever, feel free to treat this to make him/her comfortable.

Many children will also get diarrhea, usually on day 2 of the illness (some may only get diarrhea). The fluid treatment for this is the opposite of vomiting - large amounts infrequently. With diarrhea, every time the gut is challenged with something to digest, large or small, a bowel movement results. So, we try to rest the gut for hours at a time, but then challenge it with a large volume of fluid. No medications are recommended for acute diarrhea, as slowing down the intestinal motility may actually make the child sicker. We do use anti-motility agents in chronic diarrhea, but that is a different entity.

The chief goal with gastroenteritis is to prevent dehydration. The signs of dehydration are: dry lips/mucous membranes, lack of production of tears with crying, lack of urination for an extended period of time, and extreme lethargy. The risk of dehydration depends on the age of the child and the severity of the vomiting and/or diarrhea, with younger children being more susceptible. This is particularly true if the child is refusing to drink. Obviously, if the child appears to be dehydrated, the Pediatrician should be notified. If the child has persistent vomiting or appears to be getting significantly dehydrated, he/she may require intravenous fluids. A recent change in the treatment of these children is administration of a potent anti-emetic, Ondansetron (Zofran). This has prevented many children from requiring intravenous fluids, but is used only in severe cases, due to potential side-effects.

Like most illnesses in children, gastroenteritis is usually viral, so antibiotics are not indicated. In fact, treating a viral gastroenteritis with an antibiotic can result in a very serious illness known as Hemolytic-Uremic Syndrome. If the diarrhea is bloody, this can indicate a bacterial etiology, and a stool culture should be considered.

I hope this discussion proves to be useful to you (actually, I hope it doesn't, meaning your children don't get sick this year - it does happen). Remember, there are links to some excellent medical websites on my website, pinnaclepediatrics.com, as well. If all else fails, don't forget the chicken soup...

Wishing you all a happy, healthy year.

Best Regards,

Scott Serbin, M.D.

P.S. – In case your kids do get sick this winter, this issue's Back Page features some puns to put a smile back on your face. Enjoy!

Punography

I tried to catch some fog, I mist.

When chemists die, they barium.

Jokes about German sausage are the wurst.

A soldier who survived mustard gas and pepper spray is now a seasoned veteran.

I know a guy who's addicted to brake fluid. He says he can stop anytime.

I stayed up all night to see where the sun went. Then it dawned on me.

This girl said she recognized me from the vegetarian club, but I'd never met herbivore.

I'm reading a book about anti-gravity. I can't put it down.

I did a theatrical performance about puns. It was a play on words.

They told me I had type A blood, but it was a Type O.

A dyslexic man walks into a bra.

PMS jokes aren't funny. Period.

Class trip to the Coca-Cola factory. I hope there's no pop quiz.

Energizer Bunny arrested: Charged with battery.

I didn't like my beard at first. Then it grew on me.

What do you call a dinosaur with an extensive vocabulary? A thesaurus.

When you get a bladder infection, urine trouble.

What does a clock do when it's hungry? It goes back four seconds.

I wondered why the baseball was getting bigger. Then it hit me!

Broken pencils are pointless.